

| | |
|-------------------------|--|
| 1. Record Nr. | UNINA9910154750803321 |
| Autore | Bushnell C. |
| Titolo | The Admissible Dual of $GL(N)$ via Compact Open Subgroups. (AM-129), Volume 129 // C. Bushnell, P. C. Kutzko |
| Pubbl/distr/stampa | Princeton, NJ : , : Princeton University Press, , [2016] ©1993 |
| ISBN | 1-4008-8249-4 |
| Descrizione fisica | 1 online resource (327 pages) : illustrations |
| Collana | Annals of Mathematics Studies ; ; 311 |
| Classificazione | SK 340 |
| Disciplina | 512/.2 |
| Soggetti | Representations of groups Nonstandard mathematical analysis |
| Lingua di pubblicazione | Inglese |
| Formato | Materiale a stampa |
| Livello bibliografico | Monografia |
| Nota di bibliografia | Includes bibliographical references and index. |
| Nota di contenuto | Frontmatter -- Contents -- Introduction -- Comments for the reader -- 1. Exactness and intertwining -- 2. The structure of simple strata -- 3. The simple characters of a simple stratum -- 4. Interlude with Hecke algebra -- 5. Simple types -- 6. Maximal types -- 7. Typical representations -- 8. Atypical representations -- References -- Index of notation and terminology |
| Sommario/riassunto | This work gives a full description of a method for analyzing the admissible complex representations of the general linear group $G = GL(N, F)$ of a non-Archimedean local field F in terms of the structure of these representations when they are restricted to certain compact open subgroups of G . The authors define a family of representations of these compact open subgroups, which they call simple types. The first example of a simple type, the "trivial type," is the trivial character of an Iwahori subgroup of G . The irreducible representations of G containing the trivial simple type are classified by the simple modules over a classical affine Hecke algebra. Via an isomorphism of Hecke algebras, this classification is transferred to the irreducible representations of G containing a given simple type. This leads to a complete classification of the irreducible smooth representations of G , including an explicit description of the supercuspidal representations as induced representations. A special feature of this work is its virtually complete reliance on algebraic methods of a ring-theoretic kind. A full and |

accessible account of these methods is given here.
