

1. Record Nr.	UNINA9910254610703321
Autore	Benedikter Niels
Titolo	Effective Evolution Equations from Quantum Dynamics // by Niels Benedikter, Marcello Porta, Benjamin Schlein
Pubbl/distr/stampa	Cham : , : Springer International Publishing : , : Imprint : Springer, , 2016
ISBN	3-319-24898-7
Edizione	[1st ed. 2016.]
Descrizione fisica	1 online resource (97 p.)
Collana	SpringerBriefs in Mathematical Physics, , 2197-1757 ; ; 7
Disciplina	530.12
Soggetti	Quantum theory Mathematical physics Quantum Physics Mathematical Physics
Lingua di pubblicazione	Inglese
Formato	Materiale a stampa
Livello bibliografico	Monografia
Note generali	Description based upon print version of record.
Nota di bibliografia	Includes bibliographical references at the end of each chapters.
Nota di contenuto	Introduction -- Mean-Field Regime for Bosonic Systems -- Coherent States Approach.-Fluctuations Around Hartree Dynamics -- The Gross-Pitaevskii Regime -- Mean-Field regime for Fermionic Systems -- Dynamics of Fermionic Quasi-Free Mixed States -- The Role of Correlations in the Gross-Pitaevskii Energy.
Sommario/riassunto	These notes investigate the time evolution of quantum systems, and in particular the rigorous derivation of effective equations approximating the many-body Schrödinger dynamics in certain physically interesting regimes. The focus is primarily on the derivation of time-dependent effective theories (non-equilibrium question) approximating many-body quantum dynamics. The book is divided into seven sections, the first of which briefly reviews the main properties of many-body quantum systems and their time evolution. Section 2 introduces the mean-field regime for bosonic systems and explains how the many-body dynamics can be approximated in this limit using the Hartree equation. Section 3 presents a method, based on the use of coherent states, for rigorously proving the convergence towards the Hartree dynamics, while the fluctuations around the Hartree equation are considered in Section 4. Section 5 focuses on a discussion of a more subtle regime, in which the many-body evolution can be approximated

by means of the nonlinear Gross-Pitaevskii equation. Section 6 addresses fermionic systems (characterized by antisymmetric wave functions); here, the fermionic mean-field regime is naturally linked with a semiclassical regime, and it is proven that the evolution of approximate Slater determinants can be approximated using the nonlinear Hartree-Fock equation. In closing, Section 7 reexamines the same fermionic mean-field regime, but with a focus on mixed quasi-free initial data approximating thermal states at positive temperature. .

---