1. Record Nr. UNINA9910300371103321 Autore Hirano Teruyuki Titolo Measurements of Spin-Orbit Angles for Transiting Systems [[electronic resourcel 1: Toward an Understanding of the Migration History of Exoplanets / / by Teruyuki Hirano Tokyo:,: Springer Japan:,: Imprint: Springer,, 2014 Pubbl/distr/stampa **ISBN** 4-431-54586-7 Edizione [1st ed. 2014.] Descrizione fisica 1 online resource (143 p.) Collana Springer Theses, Recognizing Outstanding Ph.D. Research, , 2190-5053 521.3 Disciplina Soggetti Space sciences Planetology Observations, Astronomical Astronomy—Observations Space Sciences (including Extraterrestrial Physics, Space Exploration) and Astronautics) Astronomy, Observations and Techniques Lingua di pubblicazione Inglese **Formato** Materiale a stampa Livello bibliografico Monografia Note generali Description based upon print version of record. Nota di bibliografia Includes bibliographical references at the end of each chapters. Nota di contenuto Introduction -- Evolution History of Extrasolar Planetary Systems --Improved Modeling of the Rossiter-McLaughlin Effect -- New Observations and Improved Analyses of the Rossiter-McLaughlin Effect -- Toward the Measurements of Spin-Orbit Relations for Small Planets -- Summary and Future Prospects. This thesis presents accurate analyses of the spin-orbit angle for many Sommario/riassunto remarkable transiting exoplanetary systems, including the first measurement of the Rossiter-McLaughlin effect for a multiple transiting system. The author presents the observational methods needed to probe the spin-orbit angle, the relation between the stellar spin axis and planetary orbital axis. Measurements of the spin-orbit angle provide us a unique and valuable opportunity to understand the origin of close-in giant exoplanets, called "hot Jupiters". The first

method introduced involves observations of the Rossiter-McLaughlin effect (RM effect). The author points out the issues with the previous

theoretical modeling of the RM effect and derives a new and improved theory. Applications of the new theory to observational data are also presented for a number of remarkable systems, and the author shows that the new theory minimizes the systematic errors by applying it to the observational data. The author also describes another method for constraining the spin-orbit angle: by combining the measurements of stellar flux variations due to dark spots on the stellar surface, with the projected stellar rotational velocity measured via spectroscopy, the spin-orbit angles "along the line-of-sight" are constrained for the transiting exoplanetary systems reported by the Kepler space telescope.