

1. Record Nr.	UNINA990001126920403321
Autore	Albert, Arthur E.
Titolo	Regression and the Moore-Penrose Pseudoinverse / Arthur Albert
Pubbl/distr/stampa	New York [etc.] : Academic Press, 1972
Descrizione fisica	xiii, 180 p. ; 24 cm
Collana	Mathematics in science and engineering ; 94
Disciplina	519
Locazione	MA1 FI1
Collocazione	C-26-(94 18-235
Lingua di pubblicazione	Inglese
Formato	Materiale a stampa
Livello bibliografico	Monografia

2. Record Nr.	UNINA9910346743203321
Autore	David S. Needleman
Titolo	Emerging Approaches for Typing, Detection, Characterization, and Traceback of Escherichia coli, 2nd Edition
Pubbl/distr/stampa	Frontiers Media SA, 2018
Descrizione fisica	1 online resource (172 p.)
Collana	Frontiers Research Topics
Soggetti	Microbiology (non-medical)
Lingua di pubblicazione	Inglese
Formato	Materiale a stampa
Livello bibliografico	Monografia
Sommario/riassunto	<p>Pathogenic Escherichia coli strains cause a large number of diseases in humans, including diarrhea, hemorrhagic colitis, hemolytic uremic syndrome, urinary tract infections, and neonatal meningitis, while in animals they cause diseases such as calf scours and mastitis in cattle, post-weaning diarrhea and edema disease in pigs, and peritonitis and airsacculitis in chickens. The different E. coli pathotypes are characterized by the presence of specific sets of virulence-related genes. Therefore, it is not surprising that pathogenic E. coli constitutes a genetically heterogeneous family of bacteria, and they are continuing to evolve. Rapid and accurate molecular methods are critically needed to detect and trace pathogenic E. coli in food and animals. They are also needed for epidemiological investigations to enhance food safety, as well as animal and human health and to minimize the size and geographical extent of outbreaks. The serotype of E. coli strains has traditionally been determined using antisera raised against the >180 different O- (somatic) and 53 H- (flagellar) antigens. However, there are many problems associated with serotyping, including: it is labor-intensive and time consuming; cross reactivity of the antisera with different serogroups occurs; antisera are available only in specialized laboratories; and many strains are non-typeable. Molecular serotyping targeting O-group-specific genes within the E. coli O-antigen gene clusters and genes that are involved in encoding for the different flagellar types offers an improved approach for determining the E.</p>

coli O- and H-groups. Furthermore, molecular serotyping can be coupled with determination of specific sets of virulence genes carried by the strain offering the possibility to determine O-group, pathotype, and the pathogenic potential simultaneously. Sequencing of the O-antigen gene clusters of all of the known O-groups of *E. coli* is now complete, and the sequences have been deposited in the GenBank database. The sequence information has revealed that some *E. coli* serogroups have identical sequences while others have point mutations or insertion sequences and type as different serogroups in serological reactions. There are also a number of other ambiguities in serotyping that need to be resolved. Furthermore, new *E. coli* O-groups are being identified. Therefore, there is an essential need to resolve these issues and to revise the *E. coli* serotype nomenclature based on these findings. There are emerging technologies that can potentially be applied for molecular serotyping and detection and characterization of *E. coli*. On a related topic, the genome sequence of thousands of *E. coli* strains have been deposited in GenBank, and this information is revealing unique markers such as CRISPR (clustered regularly interspaced short palindromic repeats) and virulence gene markers that could be used to identify *E. coli* pathotypes. Whole genome sequencing now provides the opportunity to study the role of horizontal gene transfer in the evolution and emergence of pathogenic *E. coli* strains. Whole genome sequencing approaches are being investigated for genotyping and outbreak investigation for regulatory and public health needs; however, there is a need for establishing bioinformatics pipelines able to handle large amounts of data as we move toward the use of genetic approaches for non-culture-based detection and characterization of *E. coli* and for outbreak investigations.
