

1. Record Nr.	UNINA9910483366103321
Autore	Haran Shai M. J
Titolo	Arithmetical investigations : representation theory, orthogonal polynomials, and quantum interpolations // Shai M.J. Haran
Pubbl/distr/stampa	Berlin, : Springer, c2008
ISBN	3-540-78379-2
Edizione	[1st ed. 2008.]
Descrizione fisica	xii, 217 p. : ill
Collana	Lecture notes in mathematics, , 0075-8434 ; ; 1941
Disciplina	511.42
Soggetti	p-adic numbers Number theory Interpolation Representations of quantum groups
Lingua di pubblicazione	Inglese
Formato	Materiale a stampa
Livello bibliografico	Monografia
Note generali	Bibliographic Level Mode of Issuance: Monograph
Nota di bibliografia	Includes bibliographical references (p. [209]-213) and index.
Nota di contenuto	Introduction: Motivations from Geometry -- Gamma and Beta Measures -- Markov Chains -- Real Beta Chain and q-Interpolation -- Ladder Structure -- q-Interpolation of Local Tate Thesis -- Pure Basis and Semi-Group -- Higher Dimensional Theory -- Real Grassmann Manifold -- p-Adic Grassmann Manifold -- q-Grassmann Manifold -- Quantum Group $U_q(\mathfrak{su}(1, 1))$ and the q-Hahn Basis.
Sommario/riassunto	In this volume the author further develops his philosophy of quantum interpolation between the real numbers and the p-adic numbers. The p-adic numbers contain the p-adic integers Z_p which are the inverse limit of the finite rings Z/pn . This gives rise to a tree, and probability measures w on Z_p correspond to Markov chains on this tree. From the tree structure one obtains special basis for the Hilbert space $L_2(Z_p, w)$. The real analogue of the p-adic integers is the interval $[-1, 1]$, and a probability measure w on it gives rise to a special basis for $L_2([-1, 1], w)$ - the orthogonal polynomials, and to a Markov chain on "finite approximations" of $[-1, 1]$. For special (gamma and beta) measures there is a "quantum" or "q-analogue" Markov chain, and a special basis, that within certain limits yield the real and the p-adic theories. This idea can be generalized variously. In representation theory, it is the quantum general linear group $GL_n(q)$ that interpolates between the p-

adic group $GL_n(\mathbb{Z}_p)$, and between its real (and complex) analogue -the orthogonal O_n (and unitary U_n) groups. There is a similar quantum interpolation between the real and p-adic Fourier transform and between the real and p-adic (local unramified part of) Tate thesis, and Weil explicit sums.
