

- | | |
|-------------------------|---|
| 1. Record Nr. | UNISA990000528680203316 |
| Autore | CRISTIANI, Antonio |
| Titolo | Guida alle nuove norme sull'usura / Antonio Cristiani |
| Pubbl/distr/stampa | Torino : Giappichelli, 1996 |
| ISBN | 88-348-6222-8 |
| Descrizione fisica | 192 p. ; 23 cm |
| Disciplina | 345.450268 |
| Soggetti | Usura - Diritto penale |
| Collocazione | XXVI.1.C 366 (IG XI 547) |
| Lingua di pubblicazione | Italiano |
| Formato | Materiale a stampa |
| Livello bibliografico | Monografia |
-
- | | |
|-------------------------|---|
| 2. Record Nr. | UNINA9910483835103321 |
| Titolo | Biological control of insect and mite pests in Iran : a review from fundamental and applied aspects / / Javad Karimi, Hossein Madadi, editors |
| Pubbl/distr/stampa | Cham, Switzerland : , : Springer, , [2021]
©2021 |
| ISBN | 3-030-63990-8 |
| Descrizione fisica | 1 online resource (622 pages) |
| Collana | Progress in Biological Control ; ; v.18 |
| Disciplina | 632.7 |
| Soggetti | Insect pests - Biological control
Control biològic de plagues
Àcars
Insectes
Llibres electrònics |
| Lingua di pubblicazione | Inglese |
| Formato | Materiale a stampa |
| Livello bibliografico | Monografia |

Intro -- Foreword -- Preface -- Contents -- Contributors -- Chapter 1: Overview: History of Agricultural Entomology and Biological Pest Control in Iran -- 1.1 Pioneer Entomologists -- 1.2 The Early Persian Books on Entomology -- 1.3 Foundation of Karaj College of Agriculture -- 1.4 The History of Biological Pest Control in Iran -- 1.5 Hayk Mirzayans Insect Museum -- 1.6 Iranian Research Institute of Plant Protection (IRIPP) -- 1.7 Department of Biological Control -- 1.8 Other Institutes -- 1.8.1 Pasteur Institute of Iran -- 1.8.2 Razi Vaccine and Serum Research Institute -- 1.8.3 Animal Science Research Institute of Iran (ASRI) -- 1.8.4 Silkworm Breeding Company and the Silkworm Research Center -- 1.9 Plant Protection Organization (PPO) -- 1.10 Societies and Congresses -- 1.10.1 The Entomological Society of Iran (ESI) -- 1.10.2 Iranian Plant Protection Congress -- 1.10.3 Iranian Acarology -- 1.11 Medical Entomology -- 1.12 Current Biological Control Plans in Iran -- References -- Part I: Predators -- Chapter 2: Lady Beetles -- Lots of Efforts but few Successes -- 2.1 Introduction, Why Lady Beetles Are so Important? -- 2.2 The Most Important Lady Beetles Species Reported from Iran -- 2.2.1 *Adalia bipunctata* (Linnaeus 1758) -- 2.2.2 *Adalia decempunctata* (Linnaeus 1758) -- 2.2.3 *Adalia fasciatopunctata revelierei* Muls -- 2.2.4 *Adalia tetraspilota* (Hope 1831) -- 2.2.5 *Anisosticta bitriangularis* Say -- 2.2.6 *Anisosticta novemdecimpunctata* (Linnaeus 1758) -- 2.2.7 *Bromus gebleri* Weise -- 2.2.8 *Bromus octosignatus* (Gebler 1830) -- 2.2.9 *Bromus undulatus* -- 2.2.10 *Calvia quattodecimguttata* (Linnaeus 1758) -- 2.2.11 *Cheilomenes sexmaculatus* (Fabricius 1781) -- 2.2.12 *Chilocorus bipustulatus* (Linnaeus 1758) -- 2.2.13 *Clitostethus arcuatus* Rossi -- 2.2.14 *Coccinella elegantula* (Weise 1980) -- 2.2.15 *Coccinella magnopunctata* Rybakow. -- 2.2.16 *Coccinella novemnotata* Herbst -- 2.2.17 *Coccinella quattodecimpustulata* (Linnaeus 1758) -- 2.2.18 *Coccinella redemita* (Weise 1895) -- 2.2.19 *Coccinella septempunctata* (Linnaeus 1758) -- 2.2.20 *Coccinella undecimpunctata* (Linnaeus 1758) -- 2.2.21 *Cryptolaemus montrouzieri* (Mulsant 1853) -- 2.2.22 *Diloponis furschi* Ahmadi and Yazdani -- 2.2.23 *Diomus rubidus* (Motschulsky 1837) -- 2.2.24 *Exochomus flavipes* (Thunberg 1781) -- 2.2.25 *Exochomus illaesticollis* Roubal -- 2.2.26 *Exochomus melanocephalus* (Zoubkoff 1833) -- 2.2.27 *Exochomus nigripennis* Erichson -- 2.2.28 *Exochomus nigromaculatus* (Goeze, 1777) -- 2.2.29 *Exochomus pubescens* (Kuster 1848) -- 2.2.30 *Exochomus quadripustulatus* (Linnaeus 1758) -- 2.2.31 *Exochomus undulatus* (Weise 1878) -- 2.2.32 *Hippodamia (Adonia) variegata* (Goeze 1777) -- 2.2.33 *Hippodamia tredecimpunctata* (Linnaeus 1758) -- 2.2.34 *Hyperaspis duvergeri* (Fursch 1985) -- 2.2.35 *Hyperaspis concolor* (Suffrian 1843) -- 2.2.36 *Hyperaspis marmottani* Fairmaire -- 2.2.37 *Hyperaspis polita* Weise -- 2.2.38 *Hyperaspis quadrimaculata* (Redtenbacher 1844) -- 2.2.39 *Hyperaspis reppensis* (Herbst 1783) -- 2.2.40 *Hyperaspis syriaco* (Weise 1885) -- 2.2.41 *Hyperaspis vinciqerrae* Capra -- 2.2.42 *Nephus arcuatus* Kapur -- 2.2.43 *Nephus bipunctatus* (Kugelann 1794) -- 2.2.44 *Nephus biguttatus* -- 2.2.45 *Nephus bisignatus etesiacus* Fursch -- 2.2.46 *Nephus fenestratus* Sahlberg -- 2.2.47 *Nephus includes* Kirsch -- 2.2.48 *Nephus nigricans* (Weise 1879) -- 2.2.49 *Nephus quadrimaculatus* (Herbst 1783) -- 2.2.50 *Nephus ulbrichi* (Fursch 1977) -- 2.2.51 *Oenopia conglobata* (Linnaeus 1758) -- 2.2.52 *Oenopia oncina* (Olivier 1808) -- 2.2.53 *Pharoscyrnus brunneosignatus* (Marder 1949) -- 2.2.54 *Pharoscyrnus ovoideus* Anthon, P. *arabicus* Anthon and P. *setulesus* Chevrolat -- 2.2.55 *Pharoscyrnus pharoides* Marseul -- 2.2.56 *Pharoscyrnus* cf. *smirnovi* (Dobzhansky 1927).

2.2.57 *Propylea quatuordecimpunctata* (Linnaeus 1758) -- 2.2.58 *Psyllobora vigintiduopunctata* (Linnaeus 1758) -- 2.2.59 *Rodolia cardinalis* (Mulsant 1850) -- 2.2.60 *Rodolia fusti* Weise -- 2.2.61 *Scymniscus biflammatus* (Motschulsky 1837) -- 2.2.62 *Scymnus argutus* Mulsant -- 2.2.63 *Scymnus araraticus* (Iablokoff-Khnzorian 1969) -- 2.2.64 *Scymnus araxicola* (Fleischer 1900) -- 2.2.65 *Scymnus apetzi* (Mulsant 1846) -- 2.2.66 *Scymnus auritus* (Thunberg 1795) -- 2.2.67 *Scymnus flavicollis* (Redtenbacher 1844) -- 2.2.68 *Scymnus frontalis* (Fabricius 1787) -- 2.2.69 *Scymnus impexus* (Mulsant, 1850) -- 2.2.70 *Scymnus levaillanti* (Mulsant 1850) -- 2.2.71 *Scymnus mongolicus* Weiser -- 2.2.72 *Scymnus nubilus* (Mulsant 1846) -- 2.2.73 *Scymnus pallipediformis* (Gunther 1958) -- 2.2.74 *Scymnus pallipes* (Mulsant 1850) -- 2.2.75 *Scymnus quadriguttatus* (Fursch and Kreissl 1967) -- 2.2.76 *Scymnus rubromaculatus* Goeze -- 2.2.77 *Scymnus schmidt* Fursch -- 2.2.78 *Scymnus subvillosus* Goeze -- 2.2.79 *Scymnus suffrianioides* (Sahlberg 1913) -- 2.2.80 *Scymnus suffrianioides apetzoides* (Capra and Fursch 1967) -- 2.2.81 *Scymnus syriacus* (Marseul 1868) -- 2.2.82 *Scymnus testaceus* (Motschulsky 1837) -- 2.2.83 *Serangium montazerii* Fursch -- 2.2.84 *Sidis* (Nephus) *biflamulatus* -- 2.2.85 *Stethorus gilvifrons* (Mulsant 1850) -- 2.2.86 *Stethorus punctillum* (Weise 1891) -- 2.2.87 *Stethorus siphonulus* (Kapur 1948) -- 2.2.88 *Sympherobius elegans* Stephens -- 2.2.89 *Synharmonia conglobata* L. -- 2.2.90 *Thea vigintiduopunctata* L. -- 2.2.91 *Tytthaspis gebleri* (Mulsant 1850) -- 2.2.92 *Vibidia duodecimguttata* (poda, 1761) -- 2.3 Life History and Demography -- 2.4 Miscellaneous Studies -- 2.5 Predator-Prey Dynamics -- 2.5.1 Functional Response of Most Important Coccinellid Species to Different Prey Types -- 2.5.2 Numerical Response.

2.6 Population Fluctuations of *Hippodamia variegata* and Other Lady Beetle Species -- 2.7 Intraguild Predation Studies -- 2.8 Applied Studies Employing Different Coccinellid Species in Greenhouse and Field Environment -- References -- Chapter 3: Biological Control of Pests by Mites in Iran -- 3.1 Introduction -- 3.2 Predatory Mites -- 3.2.1 Order Mesostigmata -- 3.2.1.1 Doryctidae -- 3.2.1.2 Parasitidae -- 3.2.1.3 Macrochelidae -- 3.2.1.4 Phytoseiidae -- 3.2.1.4.1 Introduced Phytoseiid Mites in Iran -- 3.2.1.4.2 Native Phytoseiid Mites -- 3.2.1.4.3 Behavioural Aspects -- 3.2.1.4.4 Life Table Parameters -- 3.2.1.4.5 The Effect of Host Plants on Predatory Mites -- 3.2.1.4.6 The Side Effects of Acaricides on Biocontrol Agents -- 3.2.1.4.7 Interactions among Biocontrol Agents -- 3.2.1.4.8 Mass Rearing -- 3.2.1.5 Blattisociidae -- 3.2.1.6 Laelapidae -- 3.2.2 Suborder Prostigmata -- 3.2.2.1 Bdellidae -- 3.2.2.2 Cunaxidae -- 3.2.2.3 Tydeidae -- 3.2.2.4 Iolinidae -- 3.2.2.5 Anystidae -- 3.2.2.6 Stigmaeidae -- 3.2.2.7 Cheyletidae -- 3.3 Parasitic Mites -- 3.3.1 Order Mesostigmata -- 3.3.1.1 Otopheidomenidae -- 3.3.2 Suborder Prostigmata -- 3.3.2.1 Eriophyiidae -- 3.3.2.2 Erythraeidae -- 3.3.2.3 Trombididae -- 3.3.2.4 Other Terrestrial Parasitengonina -- 3.3.2.5 Aquatic Parasitengonina (Hydrachnidae) -- 3.3.2.6 Trochometrididae -- 3.3.2.7 Pyemotidae -- 3.3.2.8 Acarophenacidae -- 3.3.2.9 Podapolipidae -- 3.4 Conclusion -- References -- Chapter 4: Applied Ecology of Some Predacious Mites in Iran -- 4.1 Introduction -- 4.2 Important Mite Pests -- 4.2.1 Two-Spotted Spider Mite -- 4.3 Important Predatory Mites -- 4.3.1 Order Mesostigmata -- 4.3.1.1 Family Phytoseiidae -- 4.3.1.1.1 *Phytoseiulus persimilis* Athias-Henriot, 1957 -- 4.3.1.1.2 *Neoseiulus* spp. Hughes, 1948 -- 4.3.1.1.3 *Typhlodromus bagdasarjani* Wainstein and Arutunjan.

4.3.1.1.4 *Amblyseius swirskii* Athias-Henriot -- 4.3.1.1.5 Other Phytoseiid Species -- 4.3.1.2 Family Tydeidae -- 4.3.1.3 Family

Ascidae -- 4.3.1.4 Family Laelapidae -- 4.3.1.5 Family Parasitidae --
4.3.2 Order Prostigmata -- 4.3.2.1 Family Trombidiidae -- 4.3.2.2
Family Cunaxidae -- 4.3.2.3 Family Stigmaeidae -- 4.3.2.4 Family
Anystidae -- 4.3.2.5 Family Cheyletidae -- 4.4 Contribution to
Research Development -- 4.5 How Can Improve the Biological Control
by Mites in Iran? -- 4.6 Conclusions -- References -- Chapter 5:
Lacewings: Research and Applied Aspects -- 5.1 Introduction: Why
Lacewings Are Important? -- 5.2 Review on Taxonomy of Neuroptera
-- 5.3 Reports on Chrysopidae Species from Iran -- 5.4 Reports on
Coniopterygidae Species from Iran -- 5.5 Reports on Hemerobiidae
Species from Iran -- 5.6 Rearing of Green Lacewings -- 5.7 Life History
of Green Lacewings -- 5.8 Research on Green Lacewings in Iran -- 5.9
Applied Studies of Lacewings for Pest Management -- 5.10 Conclusion
-- References -- Part II: Parasitoids -- Chapter 6: Egg Parasitoids:
Chalcidoidea with Particular Emphasis on Trichogrammatidae -- 6.1
Introduction -- 6.2 Family Trichogrammatidae Foerster, 1856 -- 6.3
Importance of Egg Mortality -- 6.4 Iranian Species and Their Target
Moths -- 6.5 Laboratory Rearing on Factitious Hosts -- 6.6 Quality
Control and Parasitoid Efficacy Studies -- 6.6.1 Longevity -- 6.6.2
Fecundity -- 6.6.3 Life Table Parameters -- 6.6.4 Functional Response
-- 6.7 Targets and Inundative Programs -- 6.7.1 Codling moth, *Cydia
pomonella* L. -- 6.7.2 Carob moth, *Spectrobates ceratoniae* Zeller --
6.7.3 Cotton Bollworm, *Helicoverpa armigera* Hubner -- 6.7.3.1
Release in Cotton Fields -- 6.7.3.2 Release in Tomato Fields -- 6.7.3.3
Release in Soybean Fields -- 6.7.4 European Corn Borer, *Ostrinia
nubilalis* (Hubner) -- 6.7.5 Rice Stemborer, *Chilo suppressalis* Walker.
6.8 Comparison Between Iran and the World Status of *Trichogramma*
Research and Application.
