Record Nr.	UNINA9910485141903321
Titolo	Meta-Topolin : a growth regulator for plant biotechnology and agriculture / / Naseem Ahmad and Miroslav Strnad (editors)
Pubbl/distr/stampa	Gateway East, Singapore : , : Springer, , [2021] ©2021
ISBN	981-15-9046-X
Edizione	[1st ed. 2021.]
Descrizione fisica	1 online resource (XIII, 339 p. 42 illus., 18 illus. in color.)
Disciplina	575.97
Soggetti	Cytokinins Plant tissue culture Plant regulators Biotecnologia vegetal Citocines Cultiu de cèl·lules i teixits vegetals Llibres electrònics
Lingua di pubblicazione	Inglese
	5
Formato	Materiale a stampa
Formato Livello bibliografico	Materiale a stampa Monografia
	Materiale a stampa

1.

	improved micropropagation, rhizogenesis and ex vitro establishment of an India Ipeca (Tylophora indica Burm f.) Chapter 15. Use of meta- topolin in somatic embryogenesis Chapter 16. Meta-topolins: in vitro responses and applications in large-scale micropropagation of horticultural crops Chapter 17. Optimization of micropropagation of some woody plants using meta-topolin Chapter 18. Biotechnological application of meta-topolin as highly active aromatic cytokinin in micropropagation of medicinal plants Chapter 19. The use of the meta-topolin in tissue culture for increasing production of secondary metabolites Chapter 20. Effects of meta-topolin on the growth, physiological and biochemical parameters in plant tissue culture Chapter 21. Establishment and management of an in vitro repository of kiwifruit (Actinidia spp.) germplasmChapter 22. New generation of inhibitors of cytokinin oxidase/dehydrogenase from Arabidopsis thaliana affects shoot/root growth and seed yield Chapter 23. Topolins and related compounds and their use in agricultural applications Chapter 24. The pharmacological activity of topolins and their ribosides.
Sommario/riassunto	Plant tissue culture (PTC) technology has gained unassailable success for its various commercial and research applications in plant sciences. Plant growth regulators (PGRs) are an essential part of any plant tissue culture intervention for propagation or modification of plants. A wide range of PGRs are available, including aromatic compounds that show cytokinin activities, promote cell division and micro-propagation, viz. kinetin, N6-benzyladenine and topolins. Topolins are naturally occurring aromatic compounds that have gained popularity as an effective alternative for other frequently used cytokinins in in vitro culture of plants. Among them, meta-topolin [6-(3- hydroxybenzlyamino) purine] is the most popular and its use in plant tissue culture has amplified swiftly. During the last few decades, there have been numerous reports highlighting the effectiveness of meta- topolin in micropropagation and alleviation of various physiological disorders, rooting and acclimatization of tissue culture raised plants