

1. Record Nr.	UNINA9910696539403321
Autore	McDonald Philip Mark
Titolo	Development of vegetation in a young ponderosa pine plantation : effect of treatment duration and time since disturbance // Philip M. McDonald and Gary O. Fiddler
Pubbl/distr/stampa	Albany, CA : , : U.S. Dept. of Agriculture, Forest Service, Pacific Southwest Research Station, , [2007]
Descrizione fisica	1 online resource (23 pages) : digital, PDF file
Collana	Research paper PSW ; ; RP-251
Altri autori (Persone)	FiddlerGary O
Soggetti	Ponderosa pine - California - Plumas National Forest - Seedlings Ponderosa pine - California - Plumas National Forest - Growth Plant succession - California - Plumas National Forest Revegetation - California - Plumas National Forest
Lingua di pubblicazione	Inglese
Formato	Materiale a stampa
Livello bibliografico	Monografia
Note generali	"June 2007."
Nota di bibliografia	Includes bibliographical references (pages 21-23).
Sommario/riassunto	The density and development of deerbrush (<i>Ceanothus integerrimus</i> Hook. & Arn.), other shrubs, forbs, graminoids, and ponderosa pine (<i>Pinus ponderosa</i> Dougl. ex Laws. var. <i>ponderosa</i>) seedlings were evaluated in a young plantation in northern California from 1988 through 1997. Treatment regimes consisted of manual release (grubbing) over 3 to 6 years and resulted in vegetation recovery times of 4 to 10 years. Revegetation was remarkably diverse, rapid, and vigorous. At the end of the study, the dominant shrub, deerbrush, had significantly lower average density and foliar cover relative to the control when treated either the first 3 years or the second 3 years, but average height was similar regardless of treatment. Average density of the 11 other shrubs was statistically higher in the control relative to other treatments, but cover and height did not differ. Density, foliar cover, and height of the forbs and graminoids did not differ among treatments at the end of the study. In the control, deerbrush plants numbered 6,100 per acre; other shrubs, 2,550 per acre; forbs, 27,300 per acre; and graminoids, 5,800 plants per acre in 1997. Foliar cover of

deerbrush at 22,700 ft² per acre was more than five times that of all other naturally established species combined and slightly exceeded that of pine seedlings in all treatments.

2. Record Nr.	UNIORUON00029333
Autore	CUBINASVILI, Giorgi
Titolo	Kartuli okromcedloba VIII-XVIII saukuneebisa : albomi istoriuli mimoxilva da anotacievi = Gruzinskoe cekannoe iskusstvo s VIII po XVIII vek : Albom istoriceskij ocerk i annotacii = Georgian repoussé work VIIIth to XVIIIth centuries : A selection of half-ton plates with annotations and an essay on the developmental history of the Art = Die Georgische Goldschmiedekunst des ... / Giorgi Cubinasvili
Pubbl/distr/stampa	Tbilisi, : Sakartvelos SSR Saxelmcipo Gamomcemloba, 1957
Descrizione fisica	4 fascicoli di ca. 20 p., 200 p. di tavole sciolte ; 61 cm
Classificazione	CAU IX E
Soggetti	METALLI - GEORGIA - CATALOGO
Lingua di pubblicazione	Georgiano Russo Inglese
Formato	Materiale a stampa
Livello bibliografico	Monografia