

1. Record Nr.	UNINA9910807220503321
Autore	Bedrossian Jacob <1984->
Titolo	Dynamics near the subcritical transition of the 3D Couette flow I : below threshold case // Jacob Bedrossian, Pierre Germain, Nader Masmoudi
Pubbl/distr/stampa	Providence, RI : , : American Mathematical Society, , [2020] ©2020
ISBN	1-4704-6251-6
Descrizione fisica	1 online resource (v, 158 pages)
Collana	Memoirs of the American Mathematical Society ; ; Number 1294
Classificazione	35B3576E0576E3076F0676F1035B4076F25
Disciplina	532.58
Soggetti	Inviscid flow Mixing Shear flow Stability Three-dimensional modeling Damping (Mechanics) Viscous flow - Mathematical models
Lingua di pubblicazione	Inglese
Formato	Materiale a stampa
Livello bibliografico	Monografia
Note generali	"July 2020, volume 266, number 1294 (fourth of 6 numbers)."
Nota di bibliografia	Includes bibliographical references and index.
Nota di contenuto	Outline of the proof -- Regularization and continuation -- High norm estimate on Q2 -- High norm estimate on Q3 -- High norm estimate on Q1/0 -- High norm estimate on Q1/[not equal] -- Coordinate system controls -- Enhanced dissipation estimates -- Sobolev estimates.
Sommario/riassunto	"We study small disturbances to the periodic, plane Couette flow in the 3D incompressible Navier-Stokes equations at high Reynolds number $Re$ . We prove that for sufficiently regular initial data of size $[\epsilon]$ [less than or equal to] $c_0 Re^{-1}$ for some universal $c_0 > 0$ , the solution is global, remains within $O(c_0)$ of the Couette flow in $L^2$ , and returns to the Couette flow as $t \rightarrow [\infty]$ . For times $t \gtrsim Re^{1/3}$ , the streamwise dependence is damped by a mixing-enhanced dissipation effect and the solution is rapidly attracted to the class of "2.5 dimensional" streamwise-independent solutions referred to as streaks. Our analysis contains perturbations that experience a transient growth of kinetic energy from $O(Re^{-1})$ to $O(c_0)$ due to the algebraic linear

instability known as the lift-up effect. Furthermore, solutions can exhibit a direct cascade of energy to small scales. The behavior is very different from the 2D Couette flow, in which stability is independent of  $Re$ , enstrophy experiences a direct cascade, and inviscid damping is dominant (resulting in a kind of inverse energy cascade). In 3D, inviscid damping will play a role on one component of the velocity, but the primary stability mechanism is the mixing-enhanced dissipation. Central to the proof is a detailed analysis of the interplay between the stabilizing effects of the mixing and enhanced dissipation and the destabilizing effects of the lift-up effect, vortex stretching, and weakly nonlinear instabilities connected to the non-normal nature of the linearization"--

---