

1. Record Nr.	UNINA9910809294503321
Titolo	Carbonate systems during the Oligocene-Miocene climatic transition // edited by Maria Mutti, Werner Piller, Christian Betzler
Pubbl/distr/stampa	Hoboken, NJ, : Blackwell Pub., 2010
ISBN	1-282-72925-X 9786612729256 1-118-39836-X 1-4443-2743-7
Edizione	[1st ed.]
Descrizione fisica	1 online resource (314 p.)
Collana	Special publication ... of the international association of sedimentologists ; ; no. 42
Altri autori (Persone)	MuttiM (Maria) PillerWerner E BetzlerChristian
Disciplina	552/.58
Soggetti	Carbonate rocks Geology, Stratigraphic - Oligocene Geology, Stratigraphic - Miocene Paleoclimatology - Oligocene Paleoclimatology - Miocene
Lingua di pubblicazione	Inglese
Formato	Materiale a stampa
Livello bibliografico	Monografia
Note generali	Description based upon print version of record.
Nota di bibliografia	Includes bibliographical references and index.
Nota di contenuto	Carbonate Systems During the Oligocene-Miocene Climatic Transition; Contents; Miocene carbonate systems: an introduction; A synthesis of Late Oligocene through Miocene deep sea temperatures as inferred from foraminiferal Mg/Ca ratios; Latitudinal trends in Cenozoic reef patterns and their relationship to climate; Carbonate grain associations: their use and environmental significance, a brief review Temperate and tropical carbonate-sedimentation episodes in the Neogene Betic basins (southern Spain) linked to climatic oscillations and changes in Atlantic-Mediterranean connections: constraints from isotopic data Facies models and geometries of the Ragusa Platform (SE Sicily, Italy) near the Serravallian-Tortonian boundary; The sensitivity of a tropical foramol-rhodalgial carbonate ramp to relative sea-level change: Miocene of the central Apennines, Italy; Facies and sequence

architecture of a tropical foramol-rhodolgal carbonate ramp: Miocene of the central Apennines (Italy)  
Facies and stratigraphic architecture of a Miocene warm-temperate to tropical fault-block carbonate platform, Sardinia (Central Mediterranean Sea) Coralline algae, oysters and echinoids - a liaison in rhodolith formation from the Burdigalian of the Latium-Abruzzi Platform (Italy); Palaeoenvironmental significance of Oligocene-Miocene coralline red algae - a review; Molluscs as a major part of subtropical shallow-water carbonate production - an example from a Middle Miocene oolite shoal (Upper Serravallian, Austria)  
Echinoderms and Oligo-Miocene carbonate systems: potential applications in sedimentology and environmental reconstruction Coral diversity and temperature: a palaeoclimatic perspective for the Oligo-Miocene of the Mediterranean region; Late Oligocene to Miocene reef formation on Kita-daito-jima, northern Philippine Sea; Carbonate production in rift basins: models for platform inception, growth and dismantling, and for shelf to basin sediment transport, Miocene Sardinia Rift Basin, Italy; Index

---

Sommario/riassunto

The Oligocene and Miocene Epochs comprise the most important phases in the Cenozoic global cooling that led from a greenhouse to an icehouse Earth. Recent major advances in the understanding and time-resolution of climate events taking place at this time, as well as the proliferation of studies on Oligocene and Miocene shallow-water/neritic carbonate systems, invite us to re-evaluate the significance of these carbonate systems in the context of changes in climate and Earth surface processes. Carbonate systems, because of a wide dependence on the ecological requirements of organisms producing

---