

1. Record Nr.	UNINA9910830661503321
Autore	Holynska Malgorzata
Titolo	Single-Molecule Magnets : Molecular Architectures and Building Blocks for Spintronics
Pubbl/distr/stampa	Newark : , : John Wiley & Sons, Incorporated, , 2019 ©2018
ISBN	3-527-80991-0 3-527-80989-9
Descrizione fisica	1 online resource (451 pages)
Disciplina	541.378
Soggetti	Molecules - Magnetic properties Magnets
Lingua di pubblicazione	Inglese
Formato	Materiale a stampa
Livello bibliografico	Monografia
Nota di contenuto	Cover -- Title Page -- Copyright -- Contents -- Editorial -- Acknowledgment -- Chapter 1 Introduction to SingleMolecule Magnets -- 1.1 What Is a SingleMolecule Magnet? -- 1.1.1 SingleChain Magnets (SCMs) -- 1.1.2 Singlelon Magnets (SIMs) -- 1.1.3 Single Toroid Magnets (STMs) -- 1.2 Historical Aspects -- 1.3 Recent Progress -- 1.3.1 SMMs Based on Actinides -- 1.3.2 Organometallic SMMs -- 1.3.3 Rational Design of SMMs -- 1.3.4 Quantum Computing -- 1.3.5 SMMs in Molecular Machines -- 1.3.6 Magnetic Refrigerants -- 1.3.7 Applications in Other Disciplines -- Acknowledgment -- References -- Chapter 2 Unique Magnetic Properties -- 2.1 Introduction -- 2.2 Basic Electromagnetic Definitions -- 2.3 Magnetostatic Energy (Magnetic Work) -- 2.4 Thermodynamic Relations -- 2.5 Definition of ac Magnetic Susceptibility -- 2.6 Representative Results -- 2.6.1 ac Susceptibility Measurements in Tris(Acetylacetonato)iron(III), (Fe(acac) <sub>3</sub> ) -- 2.6.2 ac Susceptibility Measurements in a OneDimensional Chain Based on Mn <sup>6</sup> Units -- 2.6.3 Spin Relaxation in a Ferromagnetically Coupled Triangular Cu <sub>3</sub> Complex -- 2.7 Technical Aspects of the ac Susceptibility Measurements -- 2.8 Intermolecular Interactions -- 2.9 Conclusions -- References -- Chapter 3 Magnetic Modeling of Single molecule Magnets -- 3.1 Introduction -- 3.2 Atoms in a Magnetic Field -- 3.2.1 Free Atoms in a Magnetic Field -- 3.2.1.1 Lande gFactor --

3.2.2 Brillouin Theory -- 3.2.2.1  $J=1/2$  Quantum Moment -- 3.2.2.2 General Quantum Case -- 3.2.3 Energy Spectrum -- 3.2.3.1 One Electron Case -- 3.2.3.2 Many Electrons Case -- 3.2.3.3 Pauli Principle - The Two Electrons Case -- 3.2.3.4 (L,S)Multiplets - The Two Electrons Case -- 3.2.3.5 (L,S)Coupling -- 3.2.4 Crystal Fields -- 3.2.5 Singleion Anisotropy -- 3.2.5.1 Heavy RareEarths Case -- 3.2.5.2 Expressions of Hcf -- 3.2.5.3 Kramer's Theorem.

3.3 Magnetic Modeling Tools -- 3.3.1 PHI v.3.0: Software for the Analysis of Anisotropic Monomeric and Exchange Coupled Polynuclear d and fBlock Complexes -- 3.3.1.1 General Theory -- 3.3.1.2 Hamiltonian Formalism for the Exchange Coupling -- 3.3.1.3 Calculation of Thermodynamic Properties -- 3.3.1.4 Irreducible Tensor Operators (ITOs) Method -- 3.3.1.5 The Isotropic Case: Stepladder Manganese(III) Inverse[9MC3]metallacrown -- 3.3.1.6 Anisotropic Exchange Coupling -- 3.3.2 Monte Carlo Simulations: The ALPS Project Release v.2.0: Open Source Software for Strongly Correlated Systems -- 3.3.2.1 Stochastic Series Expansion (SSE) Quantum Monte Carlo Algorithm -- 3.3.2.2 Local vs Nonlocal Updates -- 3.3.2.3 Thermalization Process -- 3.3.2.4 ALPS Project: Definition of Input Files -- 3.3.2.5 The Case of Heterometallic Molecular Wheels -- 3.3.2.6 The Case of Highnuclearity Copper Cages: Tricorne Cu<sub>21</sub> and Saddlelike Cyclic Cu<sub>16</sub> -- 3.3.2.7 Other Examples. The Case of a MnIII<sub>6</sub>MnII<sub>6</sub> Molecular Wheel -- References -- Chapter 4 Insight into Magnetic and Electronic Properties Through HFEP R Studies -- 4.1 Introduction: Magnetic vs Electronic Properties of Transition Metal Ions Including SMMs and SIMs -- 4.2 Basics of HFEP R as Applied to SIMs and Other Transition Metal Complexes -- 4.2.1 Spin Hamiltonian -- 4.2.2 Methodology of Extracting ZFS and g Information from HFEP R Spectra -- 4.3 Applicability of HFEP R to Investigating SMMs and SIMs -- 4.3.1 Polynuclear Clusters -- 4.3.2 Dimers -- 4.3.3 Mononuclear Complexes -- 4.3.4 Limitations to HFEP R -- 4.3.5 Techniques Alternative to HFEP R -- 4.4 Interplay Between Spin Hamiltonian Parameters and Crystal/LigandField Parameters. From Simple Ligand Field to Sophisticated Quantum Chemical Calculations -- 4.4.1 Recapitulation -- Acknowledgment -- References -- Chapter 5 Other Techniques to Study SingleMolecule Magnets.

5.1 Introduction -- 5.2 The Mossbauer Effect -- 5.3 The Basic Principles of Mossbauer Spectroscopy -- 5.4 Hyperfine Interactions -- 5.4.1 The Isomer Shift -- 5.4.2 Quadrupole Splitting -- 5.4.3 Magnetic Hyperfine Interactions -- 5.4.4 General Remarks -- 5.5 Relaxation Phenomena and Dynamics -- 5.5.1 MixedValence Systems -- 5.6 Application of Mossbauer Spectroscopy to SingleMolecule Magnets -- 5.6.1 [FeIII<sub>8</sub>O<sub>2</sub>(OH)<sub>12</sub>(tacn)<sub>6</sub>Br<sub>89</sub>H<sub>2</sub>O -- 5.6.2 (pyrH)<sub>5</sub>[FeIII<sub>13</sub>O<sub>4</sub>F<sub>24</sub>(OMe)<sub>12</sub>4H<sub>2</sub>OMeOH -- 5.6.3 [FeIII<sub>11</sub>O<sub>7</sub>(dea)<sub>3</sub>(piv)<sub>12</sub>]Cl<sub>5</sub>MeCN -- 5.6.4 [HFeIII<sub>19</sub>O<sub>14</sub>(OEt)<sub>30</sub>] -- 5.6.5 [FeIII<sub>4</sub>(OMe)<sub>6</sub>(dpm)<sub>6</sub>] -- 5.6.6 {FeIII[FeIII(L<sub>1</sub>)<sub>2</sub>]<sub>3</sub>} -- 5.6.7 [FeII<sub>2</sub>(acpyentO)(NCO)<sub>3</sub>] -- 5.6.8 [FeII<sub>9</sub>(X)<sub>2</sub>(O<sub>2</sub>CMe)<sub>8</sub>{(2py)<sub>2</sub>CO<sub>2</sub>]<sub>4</sub>] (X = N<sub>3</sub>, NCO, OH) -- 5.6.9 [FeII<sub>7</sub>(OMe)<sub>6</sub>(Hbmsae)<sub>6</sub>]Cl<sub>26</sub>H<sub>2</sub>O -- 5.6.10 [FeIIFeIII(L)(O<sub>2</sub>CMe)<sub>2</sub>](ClO<sub>4</sub>) -- 5.6.11 [(Me<sub>3</sub>TPyA)<sub>2</sub>FeII<sub>2</sub>(L)](BARF<sub>4</sub>)<sub>2</sub> and [(Me<sub>3</sub>TPyA)<sub>2</sub>FeII/III<sub>2</sub>(L)](BARF<sub>4</sub>)<sub>3</sub>CH<sub>2</sub>Cl<sub>2</sub> -- 5.6.12 [(18C6)K(thf)<sub>2</sub>][(tbsL)Fe<sub>3</sub>] and [(crypt222)K][(tbsL)Fe<sub>3</sub>] -- 5.7 Application of Mossbauer Spectroscopy to Singleion Magnets -- 5.7.1 [M(solvn)][(tpaR)FeII] -- 5.7.2 [K(crypr222)][Fe{C(SiMe<sub>3</sub>)<sub>3</sub>]<sub>2</sub>] -- 5.7.3 [FeII{C(SiMe<sub>3</sub>)<sub>3</sub>]<sub>2</sub>] -- 5.7.4 [FeII{N(SiMe<sub>3</sub>)(Dipp)}<sub>2</sub>] -- 5.7.5 [FeII{OC<sub>6</sub>H<sub>3</sub>2,6(C<sub>6</sub>H<sub>3</sub>iPr<sub>2</sub>)<sub>2</sub>]<sub>2</sub>] -- 5.7.6 [Fe(cAAC)<sub>2</sub>Cl] -- 5.7.7 [Fe(cAAC)<sub>2</sub>][B(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>] -- 5.7.8 [K(L)][Fe{N(SiMe<sub>3</sub>)<sub>3</sub>]<sub>2</sub>] -- 5.7.9 [FeII(Eind)<sub>2</sub>] -- 5.8 Application of Mossbauer Spectroscopy to Fe/4f SingleMolecule Magnets -- 5.8.1 [FeIII<sub>4</sub>DyIII<sub>4</sub>(teaH)<sub>8</sub>(N<sub>3</sub>)<sub>8</sub>(H<sub>2</sub>O)]H<sub>2</sub>O 4MeCN -- 5.8.2 [FeIII<sub>2</sub>LnIII<sub>2</sub>(OH)<sub>2</sub>(teaH)<sub>2</sub>(O<sub>2</sub>CCPh)<sub>6</sub>]<sub>3</sub>MeCN

(LnIII = CeIII to YbIII) -- 5.8.3 [FeIII4LnIII2(teaH)4(N3)7(piv)3]  
(LnIII = YIII, GdIII, TbIII, DyIII, HoIII, ErIII) -- 5.8.4 [FeIII4DyIII2(OH)2(n  
bdea)4(C6H5CO2)8]MeCN -- 5.8.5 [FeIII4DyIII2(OH)2(nbdea)4((CH3  
3CCO2)6(N3)2)3MeCN -- 5.8.6 [Fe7Dy3( $\mu$ 4O)2( $\mu$ 3OH)2(mdea)7( $\mu$   
benzoate)4(N3)6]2H2O7MeOH -- 5.8.7 [Fe4Dy2( $\mu$ 4O)2(NO3)2(piv)6  
(Hedte)2]4MeCNC6H5OH.  
5.8.8 [FeIII2Dy2( $\mu$ 3OH)2(teg)2(N3)2(C6H5CO2)4] -- 5.8.9 [FeIII2Dy2  
( $\mu$ 3OH)2(pmide)2(pMeC6H5CO2)6] -- 5.8.10 [FeIII2DyIII2(OH)2(L1)2  
(HL2)2(NO3)4(H2O)1.5(MeOH)0.5]6MeCN -- 5.8.11 [FeIII2Ln2(H2L)4  
(NO3)2](ClO4)22MeOH2H2O (Ln = GdIII, DyIII, TbIII) -- 5.8.12  
[FeIII3Ln( $\mu$ 3O)2(CCl3CO2)8(H2O)(thf)3]x(thf)y(heptane) (LnIII = CeIII  
HoIII, LuIII, YIII) -- 5.8.13 [FeII2Dy(L)2(H2O)](ClO4)22H2O -- 5.9  
Application of Mossbauer Spectroscopy to CyanideBridged Complexes  
-- 5.10 Other Spectroscopic Techniques Used to Study IronBased  
SMMs -- 5.11 Conclusions -- References -- Chapter 6 Synthesis and  
Chemistry of Singlemolecule Magnets -- 6.1 General Introduction for  
the Synthesis of SMMs and SIMsOrganization of the Chapter -- 6.2  
Synthetic Aspects for Polynuclear 3d Metal SMMs -- 6.2.1 Approaches  
Using Simple 3d Metal Starting Materials -- 6.2.2 Approaches Using  
Preformed Coordination Clusters or SMMs as Starting  
Materials - Retention of Nuclearity -- 6.2.3 Approaches Using  
Preformed Coordination Clusters or SMMs as Starting  
Materials - Change of Nuclearity -- 6.3 Synthetic Aspects for Dinuclear  
and Polynuclear 4f Metal Complexes with SMM Properties -- 6.3.1 O  
Bridged Groups -- 6.3.2 Chlorido Bridges -- 6.3.3 Monoatomic and  
Polyatomic Nbased Ligands -- 6.3.4 Sulfurbridged SMMs -- 6.3.5  
Organometallic Bridges -- 6.3.6 Radicalbridged Lanthanide(III) SMMs  
-- 6.3.7 Multidecker Lanthanide(III)Phthalocyanine SMMs -- 6.4  
Synthetic Aspects for Dinuclear and Polynuclear Actinide SMMs -- 6.5  
Synthesis of 3d/4f, 3d/5f, 4f/5fMetal and Other Heterometallic SMMs  
-- 6.5.1 3d/4fMetal SMMs -- 6.5.2 3d/5fMetal SMMs -- 6.5.3 4f/5f  
Metal Clusters and SMMs -- 6.5.4 Other Heterometallic SMMs - the  
Synthetic Utility of the Cyano Ligand -- 6.6 Synthesis of 3d Metal SIMs  
-- 6.7 Synthetic Methodology for 4f Metal SIMs -- 6.7.1  
Phthalocyaninebased 4f Metal SIMs.  
6.7.2 Nonphthalocyanine 4f Metal SIMs -- 6.8 Synthetic Routes for 5f  
Metal SIMs -- 6.9 Concluding Comments in Brief-Prognosis for the  
Future -- References -- Chapter 7 Breakthrough in Radicalbridged  
Singlemolecule Magnets -- 7.1 General Information About Organic  
Radicals and Their Magnetic Properties -- 7.2 3d Metal Radical SMMs  
-- 7.2.1 Nitroxide Radical SMMs -- 7.2.2 Carbene Radical SMMs --  
7.2.3 Benzosemiquinonoid and Nindigo Radical SMMs -- 7.3 4f Metal  
Radical SMMs -- 7.3.1 Phthalocyanine Radical SMMs -- 7.3.2 Nitroxide  
Radical SMMs -- 7.3.3 N23 Radical SMMs -- 7.3.4 Other 4f Radical  
SMMs -- 7.4 3d-4f Metal Radical SMMs -- 7.5 5f Metal Radical SMMs  
-- 7.6 Conclusions -- References -- Chapter 8 Assembly of  
Polynuclear Singlemolecule Magnets -- 8.1 Introduction -- 8.2 History  
-- 8.3 Topological Methods in Crystal Chemistry and Coordination  
Chemistry -- 8.3.1 General Overview of ToposPro -- 8.3.2 Example of  
the ToposPro Analysis of Polynuclear Coordination Clusters -- 8.4  
Polynuclear Coordination Clusters Assembly and Topology -- 8.5 3d-4f  
PCCs -- 8.5.1 Synthetic Approach for 3d-4f PCCs -- 8.5.2 3d-4f SMMs  
PCCs -- 8.6 Assembly Examples and Graph Comparison -- 8.7  
Targeting for New Topologies -- 8.8 Synthetic Aspects in Recent  
Examples -- 8.9 Perspective -- References -- Annexure -- Chapter 1  
-- Chapter 2 -- Chapter 3 -- Chapter 4 -- Chapter 5 -- Chapter 6 --  
Chapter 7 -- Chapter 8 -- Hints/Solutions -- About the Authors --  
Index -- EULA.

---

"Concise overview of synthesis and characterization of single molecule magnets. Molecular magnetism is explored as an alternative to conventional solid-state magnetism as the basis for ultrahigh-density memory materials with extremely fast processing speeds. In particular single-molecule magnets (SMM) are in the focus of current research, both because of their intrinsic magnetization properties, as well as because of their potential use in molecular spintronic devices. Single-Molecule Magnets: Molecular Architectures and Building Blocks for Spintronics starts with a general introduction to single-molecule magnets, which helps readers to understand the evolution of the field and its future. The following chapters deal with the current synthetic methods leading to SMMs, their magnetic properties and their characterization by methods such as high-field electron paramagnetic resonance, paramagnetic nuclear magnetic resonance, and magnetic circular dichroism. The book closes with an overview of radical-bridged SMMs, which have shown application potential as building blocks for high-density memories. Covers a hot topic - single-molecule magnetism is one of the fastest growing research fields in inorganic chemistry and materials science ; Provides researchers and newcomers to the field with a solid foundation for their further work. Single-Molecule Magnets: Molecular Architectures and Building Blocks for Spintronics will appeal to inorganic chemists, materials scientists, molecular physicists, and electronics engineers interested in the rapidly growing field of study."--Page 4 of cover

---