Record Nr. UNISA996320245903316 **Titolo** Treatment of tuberculosis [[electronic resource]]: guidelines Geneva,: World Health Organization, 2010 Pubbl/distr/stampa **ISBN** 1-282-76081-5 9786612760815 92-4-068535-9 Edizione [4th ed.] Descrizione fisica 1 online resource (159 p.) Disciplina 616.995061 Soggetti Antitubercular agents Tuberculosis - Chemotherapy Patient compliance Lingua di pubblicazione Inglese **Formato** Materiale a stampa Livello bibliografico Monografia Previous editions had different title: "Treatment of tuberculosis: Note generali guidelines for national programmes". "WHO/HTM/TB/2009.420." "World Health Organization, Stop TB Dept." Includes bibliographical references. Nota di bibliografia COVER; COPYRIGHT; Contents; Abbreviations; Acknowledgements; Nota di contenuto Foreword; Executive summary; Introduction; Case definitions; Standard treatment regimens; Monitoring during treatment; Co-management of HIV and active TB disease: Supervision and patient support: Treatment of drug-resistant tuberculosis; Treatment of extrapulmonary TB and of TB in special situations; Annexes; Essential first-line antituberculosis drugs; Summary of evidence and considerations underlying the recommendations; TB treatment outcomes; Suggestions for future research Sommario/riassunto The World Health Organization's Stop TB Department has prepared this fourth edition of Treatment of Tuberculosis: Guidelines, adhering fully to the new WHO process for evidence-based guidelines. Several important recommendations are being promoted in this new edition. First, the recommendation to discontinue the regimen based on just 2 months of rifampicin (2HRZE/6HE) and change to the regimen based on

a full 6 months of rifampicin (2HRZE/4HR) will reduce the number of relapses and failures. This will alleviate patient suffering resulting from

a second episode of tuberculosis (TB) and conserve